

PrEP Risk Assessment Form

Risk Assessment – Men who have sex with men

| HIRI-MSM Risk Index* | | | |
|---|--|---------------------|--------------|
| 1 | How old are you today (yrs)? | <18 years | score 0 |
| | | 18–28 years | score 8 |
| | | 29–40 years | score 5 |
| | | 41–48 years | score 2 |
| | | ≥49 years | score 0 |
| 2 | How many men have you had sex with in the last 6 months? | >10 male partners | score 7 |
| | | 6–10 male partners | score 4 |
| | | 0–5 male partners | score 0 |
| 3 | In the last 6 months, how many times did you have receptive anal sex (you were the bottom) with a man? | 1 or more times | score 10 |
| | | 0 times | score 0 |
| 4 | How many of your male sex partners were HIV positive? | >1 positive partner | score 8 |
| | | 1 positive partner | score 4 |
| | | <1 positive partner | score 0 |
| 5 | In the last 6 months, how many times did you have insertive anal sex (you were the top) with a man who was HIV positive? | 5 or more times | score 6 |
| | | 0 times | score 0 |
| 6 | In the last 6 months, have you used methamphetamines such as crystal or speed? | Yes | score 5 |
| | | No | score 0 |
| 7 | In the last 6 months, have you used poppers (amyl nitrate)? | Yes | score 3 |
| | | No | score 0 |
| Add down entries in right column to calculate total score | | | Total score† |
| <p>*To identify sexually active MSM in their practice, we recommend clinicians ask all their male patients a routine question: "In the past (time) have you had sex? (if yes), with men, women, or both?"</p> <p>†If score is 10 or greater, evaluate for PrEP or other intensive HIV prevention services; If score is 9 or less, provide indicated standard HIV prevention services.</p> | | | |

Risk Assessment – Heterosexual Men & Women

In the past 6 months:

- How many men/women have you had sex with?
- How many times did you have vaginal or anal sex when neither you nor your partner wore a condom?
- How many of your sex partners were HIV-positive?
- (*if any positive*) With these HIV-positive partners, how many times did you have vaginal or anal sex without a condom?

Risk Assessment Tool – Intravenous Drug User

- When did you last inject unprescribed drugs?
- In the past 6 months, have you injected by using needles, syringes, or other drug preparation equipment that had already been used by another person?
- In the past 6 months, have you been in a methadone or other medication-based drug treatment program?